

Crosstab

			kejadian pendek		Total
			Stunting	Tidak Stunting	
kualitas sumber air	Sumber air bersih baik	Count	26	28	54
		% within kualitas sumber air	48.1%	51.9%	100.0%
		% within kejadian pendek	68.4%	73.7%	71.1%
	Sumber air tidak bersih	% of Total	34.2%	36.8%	71.1%
		Count	12	10	22
		% within kualitas sumber air	54.5%	45.5%	100.0%
		% within kejadian pendek	31.6%	26.3%	28.9%
	Total	% of Total	15.8%	13.2%	28.9%
		Count	38	38	76
		% within kualitas sumber air	50.0%	50.0%	100.0%
		% within kejadian pendek	100.0%	100.0%	100.0%
		% of Total	50.0%	50.0%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.256 ^a	1	.613	.801	.400
Continuity Correction ^b	.064	1	.800		
Likelihood Ratio	.256	1	.613		
Fisher's Exact Test					
Linear-by-Linear Association	.253	1	.615		
N of Valid Cases	76				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 11.00.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for kualitas sumber air (Sumber air bersih baik / Sumber air tidak bersih)	.774	.286	2.092
For cohort kejadian pendek = Stunting	.883	.551	1.414
For cohort kejadian pendek = Tidak Stunting	1.141	.675	1.928
N of Valid Cases	76		

sumber air minum * kejadian pendek Crosstabulation

		kejadian pendek		Total
		Stunting	Tidak Stunting	
sumber air minum	Count	18	9	27
	tidak baik			
	% within sumber air minum	66.7%	33.3%	100.0%
	% within kejadian pendek	47.4%	23.7%	35.5%
	% of Total	23.7%	11.8%	35.5%
	Count	20	29	49
	baik			
	% within sumber air minum	40.8%	59.2%	100.0%
	% within kejadian pendek	52.6%	76.3%	64.5%
Total	% of Total	26.3%	38.2%	64.5%
	Count	38	38	76
	% within sumber air minum	50.0%	50.0%	100.0%
	% within kejadian pendek	100.0%	100.0%	100.0%
	% of Total	50.0%	50.0%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	4.653 ^a	1	.031	.054	.027
Continuity Correction ^b	3.676	1	.055		
Likelihood Ratio	4.721	1	.030		
Fisher's Exact Test					
Linear-by-Linear Association	4.592	1	.032		
N of Valid Cases	76				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 13,50.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for sumber air minum (tidak baik / baik)	2.900	1.086	7.744
For cohort kejadian pendek = Stunting	1.633	1.063	2.511
For cohort kejadian pendek = Tidak Stunting	.563	.315	1.008
N of Valid Cases	76		

Sarana SPAL * kejadian pendek Crosstabulation

		kejadian pendek		Total
		Stunting	Tidak Stunting	
Sarana SPAL	Count	30	20	50
	Tidak baik			
	% within Sarana SPAL	60.0%	40.0%	100.0%
	% within kejadian pendek	78.9%	52.6%	65.8%
	% of Total	39.5%	26.3%	65.8%
	Count	8	18	26
	baik			
	% within Sarana SPAL	30.8%	69.2%	100.0%
	% within kejadian pendek	21.1%	47.4%	34.2%
	% of Total	10.5%	23.7%	34.2%
Total	Count	38	38	76
	% within Sarana SPAL	50.0%	50.0%	100.0%
	% within kejadian pendek	100.0%	100.0%	100.0%
	% of Total	50.0%	50.0%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	5.846 ^a	1	.016	.029	.014
Continuity Correction ^b	4.735	1	.030		
Likelihood Ratio	5.961	1	.015		
Fisher's Exact Test					
Linear-by-Linear Association	5.769	1	.016		
N of Valid Cases	76				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 13,00.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Sarana SPAL (Tidak baik/ baik)	3.375	1.233	9.237
For cohort kejadian pendek = Stunting	1.950	1.050	3.623
For cohort kejadian pendek = Tidak Stunting	.578	.378	.884
N of Valid Cases	76		

jamban keluarga * kejadian pendek Crosstabulation

		kejadian pendek		Total
		Stunting	Tidak Stunting	
jamban keluarga	Count	29	19	48
	% within jamban keluarga	60.4%	39.6%	100.0%
	% within kejadian pendek	76.3%	50.0%	63.2%
	% of Total	38.2%	25.0%	63.2%
	Count	9	19	28
	% within jamban keluarga	32.1%	67.9%	100.0%
	% within kejadian pendek	23.7%	50.0%	36.8%
	% of Total	11.8%	25.0%	36.8%
	Count	38	38	76
Total	% within jamban keluarga	50.0%	50.0%	100.0%
	% within kejadian pendek	100.0%	100.0%	100.0%
	% of Total	50.0%	50.0%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	5.655 ^a	1	.017		
Continuity Correction ^b	4.580	1	.032		
Likelihood Ratio	5.750	1	.016		
Fisher's Exact Test				.031	.016
Linear-by-Linear Association	5.580	1	.018		
N of Valid Cases	76				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 14,00.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for jamban keluarga (Tidak Sehat/ Sehat)	3.222	1.207	8.600
For cohort kejadian pendek = Stunting	1.880	1.047	3.373
For cohort kejadian pendek = Tidak Stunting	.583	.378	.899
N of Valid Cases	76		

Penanganan kotoran baduta * kejadian pendek Crosstabulation

			kejadian pendek		Total
			Stunting	Tidak Stunting	
Penanganan kotoran baduta	Tidak Baik	Count	27	17	44
		% within Penanganan kotoran baduta	61.4%	38.6%	100.0%
		% within kejadian pendek	71.1%	44.7%	57.9%
		% of Total	35.5%	22.4%	57.9%
	Baik	Count	11	21	32
		% within Penanganan kotoran baduta	34.4%	65.6%	100.0%
		% within kejadian pendek	28.9%	55.3%	42.1%
		% of Total	14.5%	27.6%	42.1%
Total	Count	38	38	76	
	% within Penanganan kotoran baduta	50.0%	50.0%	100.0%	
	% within kejadian pendek	100.0%	100.0%	100.0%	
	% of Total	50.0%	50.0%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	5.398 ^a	1	.020	.036	.018
Continuity Correction ^b	4.372	1	.037		
Likelihood Ratio	5.471	1	.019		
Fisher's Exact Test					
Linear-by-Linear Association	5.327	1	.021		
N of Valid Cases	76				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 16,00.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Penanganan kotoran baduta (Tidak Baik / Baik)	3.032	1.174	7.831
For cohort kejadian pendek = Stunting	1.785	1.048	3.042
For cohort kejadian pendek = Tidak Stunting	.589	.376	.922
N of Valid Cases	76		

Pengelolaan sampah rumah tangga * kejadian pendek Crosstabulation

		kejadian pendek		
		Stunting	Tidak Stunting	
Pengelolaan sampah rumah tangga	Tidak Baik	Count	27	17
		% within Pengelolaan sampah rumah tangga	61.4%	38.6%
		% within kejadian pendek	71.1%	44.7%
		% of Total	35.5%	22.4%
	baik	Count	11	21
		% within Pengelolaan sampah rumah tangga	34.4%	65.6%
		% within kejadian pendek	28.9%	55.3%
		% of Total	14.5%	27.6%
Total	Count	38	38	
	% within Pengelolaan sampah rumah tangga	50.0%	50.0%	
	% within kejadian pendek	100.0%	100.0%	
	% of Total	50.0%	50.0%	

Chi-Square Tests					
	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	5.398 ^a	1	.020	.036	.018
Continuity Correction ^b	4.372	1	.037		
Likelihood Ratio	5.471	1	.019		
Fisher's Exact Test					
Linear-by-Linear Association	5.327	1	.021		
N of Valid Cases	76				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 16,00.

b. Computed only for a 2x2 table

Risk Estimate			
	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Pengelolaan sampah rumah tangga (Tidak Baik/ baik)	3.032	1.174	7.831
For cohort kejadian pendek = Stunting	1.785	1.048	3.042
For cohort kejadian pendek = Tidak Stunting	.589	.376	.922
N of Valid Cases	76		