

- Turnitin

Contangion ACI

 task 3

Document Details

Submission ID

trn:oid::3618:124665044

Submission Date

Dec 17, 2025, 11:45 AM GMT+7

Download Date

Dec 17, 2025, 11:50 AM GMT+7

File Name

Contangion ACI.docx

File Size

1.2 MB

10 Pages

3,746 Words

24,648 Characters

15% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

Match Groups

-  **40 Not Cited or Quoted 9%**
Matches with neither in-text citation nor quotation marks
-  **0 Missing Quotations 0%**
Matches that are still very similar to source material
-  **6 Missing Citation 1%**
Matches that have quotation marks, but no in-text citation
-  **0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 9%  Internet sources
- 4%  Publications
- 0%  Submitted works (Student Papers)

Match Groups

- 40 Not Cited or Quoted 9%**
Matches with neither in-text citation nor quotation marks
- 0 Missing Quotations 0%**
Matches that are still very similar to source material
- 6 Missing Citation 1%**
Matches that have quotation marks, but no in-text citation
- 0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 9% Internet sources
- 4% Publications
- 0% Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Internet	jurnal.uinsu.ac.id	2%
2	Internet	jurnalku.org	<1%
3	Internet	www.mdpi.com	<1%
4	Internet	e-journal.unair.ac.id	<1%
5	Internet	journal.uinsgd.ac.id	<1%
6	Publication	Cindy Chevalley-Gerber, Isabelle Carrard, Marielle Schmied, Raphaël Hammer. "F...	<1%
7	Internet	prin.or.id	<1%
8	Internet	projekter.aau.dk	<1%
9	Internet	www.scilit.net	<1%
10	Internet	ejournal.lucp.net	<1%

11	Publication	Chagit Peles, Netalie Shloim, Mary C.J. Rudolf. "Nutritional Nesting of First-Time E...	<1%
12	Publication	Wenting Jiang, Lijun Chen. "Analysis of the factors and moderating role of self-car...	<1%
13	Internet	advancesinresearch.id	<1%
14	Internet	archive.org	<1%
15	Internet	journal.stikeshangtuah-sby.ac.id	<1%
16	Internet	online-journal.unja.ac.id	<1%
17	Internet	proceeding.unisayogya.ac.id	<1%
18	Internet	ejurnal.poltekkes-tjk.ac.id	<1%
19	Internet	library.upol.cz	<1%
20	Internet	orca.cardiff.ac.uk	<1%
21	Internet	theses.hal.science	<1%
22	Internet	jurnal.poltekkeskupang.ac.id	<1%
23	Internet	ejournal.unuja.ac.id	<1%
24	Internet	rcastoragev2.blob.core.windows.net	<1%

25	Publication	Luis Gustavo Saboia Ponte, Suliene França Ribeiro, Adriane Elisabete Costa Antun...	<1%
26	Internet	openresearch.lsbu.ac.uk	<1%
27	Internet	ouci.dntb.gov.ua	<1%
28	Internet	www.davaodoctors.edu.ph	<1%
29	Internet	www.scielo.br	<1%



The Relationship between Self-Efficacy in Choosing Nutritious Foods and the Daily Consumption Practices of Pregnant Women

Asri Masitha Arsyati¹, Nhadira Mhesticia², Agustina³

¹ Department of Public Health of Ibn Khaldun University, Bogor

² Faculty of Mathematics and Natural Sciences of Pakuan University, Bogor
Departement of Nursing of Poltekkes Bandung

Correspondence Email: asri.masitha@uika-bogor.ac.id

Track Record Article

Accepted:

Published:

Abstract

Adequate consumption of nutritious food by pregnant women is a crucial factor in supporting maternal health and fetal growth and development. However, various factors such as knowledge, self-efficacy, and social support can influence the daily consumption behavior of pregnant women. The purpose of this study is to determine the determinants associated with adequate daily consumption in pregnant women. An observational analytical study with a cross-sectional design was conducted on 231 pregnant women. The relationship between variables was analyzed using the Chi-Square statistical test with a p-value <0.05 as the significance threshold. Mothers' self-efficacy in choosing nutritious foods is significantly related to daily consumption adequacy ($p = 0.000$; OR = 0.319). Nutritional knowledge also shows a significant relationship ($p = 0.016$; OR = 2.043). Family social support has a significant effect on the daily consumption of pregnant women ($p = 0.008$; OR = 2.780). Meanwhile, food access ($p = 0.978$; OR = 0.976), antenatal nutrition counseling ($p = 0.148$; OR = 0.648), access to pregnancy nutrition information media ($p = 0.116$; OR = 2.872), and cultural/food taboo practices ($p = 0.246$; OR = 0.644) do not show a significant relationship. Self-efficacy, nutrition knowledge, and family social support are factors that significantly influence the adequacy of daily consumption among pregnant women. Efforts to improve nutrition education and strengthen family support should be the focus of interventions aimed at encouraging healthy eating behaviors among pregnant women.

Keyword: Food consumption of pregnant women, self-efficacy, nutritional knowledge, family social support, nutritional determinants of pregnant women.

INTRODUCTION

Pregnancy in a woman's life is a critical period, during which nutritional needs increase significantly to support fetal growth and maternal health (Christy & Simanjuntak, 2023; Kabahenda & Stoecker, 2024). Globally, micronutrient and energy deficiencies during pregnancy are associated with risks of fetal growth disorders, low birth weight, and other obstetric complications (Olloqui-Mundet et al., 2024). A study conducted in various countries, shows that the consumption patterns of pregnant women often do not meet nutritional recommendations during pregnancy (Fernández-Gómez et al., 2020).

Author names dkk / Scientific Periodical of Public Health and Coastal 4(2),2022 , halaman 87-99 (10pt, all caps, Cambria, Normal)

At the national level, in Indonesia, nutritional fulfillment in pregnant women remains a challenge and a special concern. For example, a study at the Kemayoran Community Health Center found that the eating habits of pregnant women are closely related to the nutritional status of mothers and the potential risk of stunting in children in the future (Mulyati et al., 2025). In addition, a study conducted at the Kuta Padang Layung health center reported that of the pregnant women surveyed, most had optimal energy and protein intake, but there were significant variations in the consumption of macronutrients such as fat and carbohydrates, as well as daily meal frequency (Nabillah et al., 2024).

This issue is important to examine because the daily consumption of pregnant women not only affects the health of the mother and the growth of the fetus, but also has social and cultural implications. Socially, malnutrition can increase the burden on public health and reduce the quality of life. Culturally, factors such as food taboos, maternal education, and nutritional awareness also shape eating patterns during pregnancy. For example, the literature notes that maternal nutritional awareness is greatly influenced by factors such as education, culture, and access to information during pregnancy (Fang et al., 2024).

Although a number of studies have explored the determinants of nutritional intake among pregnant women (e.g., socioeconomic status, education, and dietary recommendations), there is a significant gap in the literature: few quantitative studies have combined analyses of social, cultural, and behavioral factors to explain variations in daily intake among pregnant women (Hanani et al., 2016). In addition, nutritional interventions during pregnancy, although showing positive results (e.g., increased micronutrient intake and reduced pregnancy complications) (Yu et al., 2024), have not been studied in depth in relation to structural factors such as income, local culture, and access to health services.

Based on this background, this study aims to analyze the factors that influence the daily consumption of pregnant women in Indonesia, focusing on the variables of self-efficacy, nutritional knowledge, food access, family social support, antenatal nutritional counseling, access to nutritional information media, and cultural customs/restrictions. Quantitatively, this study will examine the relationship between these factors and the daily consumption of pregnant women. The theoretical contribution of this study is to enrich the framework of determinants of maternal nutrition by incorporating cultural factors and self-efficacy into a quantitative model, while the practical contribution includes policy recommendations and more targeted nutritional intervention programs for pregnant women in the community.

METHODS

1

9

14

This study used a cross-sectional design and was conducted in the working area of maternal and child health facilities in Bogor city. The location was chosen based on the consideration that there was a well-run nutritional monitoring program for pregnant women, making it easier to access data and respondents. The study took place in April. The population in this study consisted of all pregnant women who were registered and actively undergoing antenatal care in the working area. The inclusion criteria included pregnant women in any trimester, willingness to be a respondent, ability to communicate well, and no health disorders that could severely affect consumption patterns. From the available population, a sample of 231 pregnant women was determined. The sampling technique used was purposive sampling. The research instrument consisted of a structured questionnaire covering sociodemographic characteristics, individual aspects (pregnant women's nutritional knowledge and self-efficacy), environmental aspects (family support), and contextual aspects (access to food, information media, and cultural/taboos). In addition, a Food Frequency Questionnaire (FFQ) form was used to measure daily food consumption. Data analysis was performed descriptively and inferentially using the Chi-square test. This study has received research ethics approval from the STIKes Hang Tuah Surabaya Research Ethics Committee Number: PE/285/II/2025/KEP/SHT.

RESULTS

Table 1 Characteristics of Respondents (n=231)

Characteristics		
	Frequency	Percentage (%)
Specific Pregnancy Conditions (e.g. Anemia/Chronic Energy Deficiency)		
Yes	95	41.1
No	136	58.9
Mother's Education		
Low	140	60.6
High	91	39.4
Household Income		
Low	200	86.6
High	31	13.4
Number of Household Members		
>3	156	67.5
≤3	75	32.5
Mother's Occupation		
Unemployed	123	53.2

Page
PAG
E
MER
GEF
ORM
AT
L

Author names dkk / Scientific Periodical of Public Health and Coastal 4(2),2022 , halaman 87-99 (10pt, all caps, Cambria, Normal)

Employed 108 46.8

Descriptive analysis shows that most respondents did not have specific pregnancy conditions such as anemia or chronic energy deficiency (58.9%), although the proportion of pregnant women with these conditions was still quite large (41.1%). In terms of education, the majority of mothers were in the low education group (60.6%), and only 39.4% had higher education. Household income was mostly in the low category (86.6%), with only 13.4% having high income. The household structure was dominated by households with more than three members (67.5%). The employment status of mothers showed a relatively balanced distribution, but slightly more mothers were unemployed (53.2%) than employed (46.8%).

Table 2 Determinants Affecting Daily Consumption of Pregnant Women

Variables	Daily Consumption of Pregnant Women						P-Value	OR
	Lack		Adequate		Total			
	n	%	n	%	n	%		
Mother's Self-Efficacy in Choosing Nutritional Foods								
Low	35	49.9	99	84.1	134	100	0.000	0.319
High	51	36.1	46	60.9	97	100		(0.183-0.555)
Nutrition Knowledge								
Inadequate	58	48.8	73	82.2	131	100	0.016	2.043
Good	28	37.2	72	62.8	100	100		(1.171-3.563)
Food Access								
Difficult	37	37.6	64	63.4	101	100	0.978	0.956
Easy	49	48.4	81	81.6	130	100		(0.558-1.637)
Family Social Support								
Less supportive	75	66.3	103	111.7	178	100	0.008	2.780
Supportive	11	19.7	42	33.3	53	100		(1.343-5.755)
Antenatal Nutrition Conseling								
Rarely	43	48.8	88	82.2	131	100	0.148	0.648
Often	43	37.2	57	62.8	100	100		(0.378-1.110)
Access to Pregnancy Nutrition Information Media								
Rarely	8	4.8	5	8.2	13	100	0.116	2.872
Often	78	81.2	140	136.8	218	100		(0.908-9.080)
Cultural Customs/Taboos								
Yes	16	20.1	38	33.9	54	100	0.246	0.648
No	70	65.9	107	111.1	177	100		(0.334-1.242)

DISCUSSION

Self-Efficacy

The results showed that pregnant women with high self-efficacy in choosing nutritious foods were more likely to consume adequate daily nutrition than women with low self-efficacy ($OR < 1, p = 0.000$). This indicates that mothers' confidence in their ability to choose nutritious foods is a protective factor for good nutrition. This phenomenon is in line with Albert Bandura's self-efficacy theory, which states that an individual's belief in their own abilities influences their intended behavior. A study by Emamipour et al., (2024) states that self-efficacy-based interventions can improve eating patterns in overweight pregnancies. In addition, a study in Korea also found that pregnant women with high self-efficacy tend to maintain a healthier diet (Kim, 2025). In the context of daily consumption by pregnant women, high self-efficacy may encourage mothers to be more active in seeking, selecting, and preparing nutritious foods, for example, prioritizing vegetables, fruits, and adequate protein, as well as overcoming obstacles such as nausea or financial constraints. Conversely, low self-efficacy can cause mothers to feel unable to change their eating patterns, making it easier for them to fall into suboptimal consumption patterns.

From the perspective of implementing nutrition programs for pregnant women, these findings underscore the importance of building psychosocial aspects such as self-confidence and empowerment in pregnant women so that they are able to choose nutritious foods independently. Strategies such as food selection skills training, nutritious menu demonstrations, and peer support can help increase self-efficacy. As a complementary theory, models such as the Health Action Process Approach (HAPA) also emphasize self-efficacy as one of the main constructs in health behavior change (Schwarzer & Hamilton, 2020).

Nutrition Knowledge for Pregnant Women

The study shows that pregnant women who have "good nutritional knowledge" also have a higher chance of adequate daily consumption ($OR > 1, p = 0.016$). This means that nutritional knowledge plays a significant role in food consumption. This is consistent with a number of studies showing that pregnant women's nutritional knowledge is positively correlated with better eating behaviors (the relationship between pregnancy nutrition knowledge, food safety, and food intake frequency) (Adylbekova et al., 2025) as well as studies in Indonesia showing that pregnant women with low nutritional knowledge tend to have less diverse consumption patterns (Angraini et al., 2023). Nutrition knowledge is not

Author names dkk / Scientific Periodical of Public Health and Coastal 4(2),2022 , halaman 87-99 (10pt, all caps, Cambria, Normal)

only about knowing what to eat, but also understanding why, when, and how to plan daily meals to meet the needs of mothers and fetuses (Nasution & Nasution, 2020; Olloqui-Mundet et al., 2024). Good knowledge enables pregnant women to recognize the benefits of nutrients such as iron, calcium, folic acid, and the negative effects of consuming inappropriate foods. In addition, knowledge also facilitates mothers in addressing consumption barriers such as nausea, cultural taboos, or economic limitations. In terms of policy, improving the nutritional knowledge of pregnant women must be integrated into antenatal care programs, nutritional counseling, and community education. Importantly, knowledge is not enough if it is not accompanied by skills and access support (Nasution et al., 2024; Engidaw et al., 2025; Olloqui-Mundet et al., 2024). The results show that despite significant knowledge, several other factors (food access, culture/taboo) do not all show a strong relationship.

Food Access

Although in this study the food access variable showed an OR = 0.956 ($p = 0.978$) and was not significant, it is still important to discuss it theoretically because the literature refers to food access as an important determinant of pregnant women's consumption. For example, a study in Singapore found that food insecurity can still hinder the quality of pregnant women's diets even though they appear to be anthropometrically normal (Lim et al., 2025). In the context of Indonesia or developing countries, food access can take the form of food availability in households, ease of purchasing nutritious food, food quality, and economic ability to purchase food. If access is limited, for example, if pregnant women live far from markets, food prices are high, or choices of nutritious food are limited, daily consumption can be disrupted. Although the results of this study were not significant, this may be because food access in the population was fairly homogeneous or lacked variability, so the effects did not appear. However, in theory, nutritional interventions for pregnant women should also include efforts to improve food access (food subsidies, strengthening local supply, food programs for pregnant women) so that knowledge and self-efficacy can be translated into actual consumption.

Family Social Support

The findings show that social support from family is significantly associated with adequate daily consumption by pregnant women (OR = 2.780; $p = 0.008$). This indicates that pregnant women who receive social support from their families are more than twice as likely to achieve adequate consumption compared to those who have less family support. These findings are consistent with international literature showing that social support during pregnancy plays an increasingly important role in the physical health and behavior of pregnant

Author names dkk / Scientific Periodical of Public Health and Coastal 4(2),2022, halaman 87-99 (10pt, all caps, Cambria, Normal)

women. For example, Al-Mutawtah et al., (2023) A systematic review shows that emotional, instrumental, and informational support from family, partners, and friends greatly affects the well-being of pregnant women.

Theoretically, social support can act as a buffer against physical and psychosocial stress during pregnancy and facilitate access, motivation, and concrete actions in nutritional behavior. In Cohen and Wills' stress buffer model, social support not only reduces the effects of stress but also increases self-efficacy and healthy behavior (Al-Mutawtah et al., 2023). In practice, supportive families might help pregnant women choose nutritious foods, prepare meals, encourage them to attend antenatal checkups, or accompany them through nutrition education. Interventions involving the pregnant woman's family and social environment are highly relevant.

Antenatal Nutrition Counseling

The variable “antenatal nutrition counseling (often vs rarely)” in this study did not show a significant relationship ($OR = 0.648$; $p = 0.148$) with daily consumption. This is somewhat surprising given that the literature shows that nutrition counseling or education during pregnancy tends to have an impact on maternal nutritional behavior. For example, a study in Indonesia shows that nutrition education can increase the knowledge of pregnant women, although this does not always translate into different pregnancy outcomes (Suryani et al., 2024).

There are several possible reasons why nutritional counseling was not significant in this study: the quality or frequency of counseling may have been inadequate, the material may have been insufficient to change daily consumption, or the measurement of daily consumption used may have been insufficiently sensitive to the effects of counseling. From a behavioral change theory perspective, simply providing information (educational outreach) is not enough if it is not followed by motivation (self-efficacy), a supportive environment (social support + access to food), and the mother's readiness for change (Iwanowicz-Palus et al., 2022; Rockliffe et al., 2021). This reiterates that nutritional interventions must be multifaceted.

Access to Pregnancy Nutrition Information Media

Although the variable of access to pregnancy nutrition information media showed an $OR = 2.872$, $p = 0.116$ (not significant). However, in the literature, access to information through media (internet, social media, leaflets) is increasingly considered important for improving the nutritional literacy of pregnant women. For example, a study by Nilawati et al., (2024) shows that video-based education can improve the nutritional knowledge and behavior

Author names dkk / Scientific Periodical of Public Health and Coastal 4(2),2022 , halaman 87-99 (10pt, all caps, Cambria, Normal)

of pregnant women. From the perspective of behavioral communication theory, the use of information media can strengthen cognitive (knowledge), affective (self-confidence), and conative (intention) factors in behavioral change models such as HAPA or Theory of Planned Behaviour (Hagger & Hamilton, 2025; Ju et al., 2025). Although not significant in this study, media access can still be considered a supporting element.

Cultural Customs/Taboos

The variable “cultural customs/taboo” also showed an OR = 0.644 and p = 0.246 (not significant), so in this study it was not proven to be a major determinant of daily consumption by pregnant women. However, it is still important to discuss. Nutritional literacy among pregnant women shows that cultural food taboos, such as not eating certain fish, eggs or vegetables, have the potential to inhibit the consumption of nutritious foods. A study in Indonesia shows that low knowledge and cultural taboos are risk factors for the nutritional consumption of pregnant women (Suryani et al., 2024). In theory, cultural factors influence eating attitudes and behaviors through social norms, risk perceptions, and traditional beliefs. Comprehensive behavior change models (e.g., HAPA) emphasize that in addition to beliefs and knowledge, contextual factors such as social and cultural norms also play an important role (Monterrosa et al., 2020; Çoker et al., 2022). Although not significant in this study, in the context of maternal nutrition programs in Indonesia and similar countries, it is very important to pay attention to cultural aspects, including culturally sensitive education and the involvement of community leaders.

CONCLUSIONS

Three variables that were found to have a significant relationship with adequate daily consumption were self-efficacy in choosing nutritious foods, nutritional knowledge, and family social support. Meanwhile, other factors such as food access, antenatal nutrition counseling, access to media information, and cultural customs/restrictions did not show a significant relationship in this study.

REFERENCE

- Adylbekova, P., Balantekin, K., Feeser, K., & Temple, J. (2025). Association among nutrition knowledge, food frequency, and food insecurity during pregnancy. *BMC Nutrition*, 11(1), 69. <https://doi.org/10.1186/s40795-025-01051-z>
- Al-Mutawtah, M., Campbell, E., Kubis, H. P., & Erjavec, M. (2023). Women’s experiences of social support during pregnancy: a qualitative systematic review. *BMC Pregnancy and Childbirth*, 23(1). <https://doi.org/10.1186/s12884-023-06089-0>

Author names dkk / Scientific Periodical of Public Health and Coastal 4(2),2022 , halaman 87-99 (10pt, all caps, Cambria, Normal)

- 1 Angraini, D. I., Sari, M. I., & Imantika, E. (2023). Nutrition Knowledge During Pregnancy in Malnutrition Pregnancy Women in Bandar Lampung City: A Qualitative Study. *The International Conference on Social Determinants of Health Research, Icsdh 2023*, 49–55. <https://doi.org/10.5220/0012898700004564>
- 9 Christy, J., & Simanjuntak, E. (2023). The Relationship Between the Nutritional Status of Pregnant Women, Infant Birth Weight and Stunting Incidence in Toddlers. *Contagion: Scientific Periodical Journal of Public Health and Coastal Health*, 5(1), 23. <https://doi.org/10.30829/contagion.v5i1.14952>
- 23 Coker, E. N., Jebb, S. A., Stewart, C., Clark, M., & Pechey, R. (2022). Perceptions of social norms around healthy and environmentally-friendly food choices: Linking the role of referent groups to behavior. *Frontiers in Psychology*, 13, 974830. <https://doi.org/10.3389/fpsyg.2022.974830>
- 20 Emamipour, F., Rajati, M., Ashtarian, H., & Rajati, F. (2024). The Effect of a Self-Efficacy-Based Dietary Intervention on Weight Gain Control in Obese and Overweight Pregnant Women: A Quasi-Experimental Pretest-Posttest Design Study. *Middle East Journal of Rehabilitation and Health Studies*, 11(4). <https://doi.org/10.5812/mejrh-144863>
- Engidaw, M. T., Lee, P., Fekadu, G., Mondal, P., & Ahmed, F. (2025). Effect of Nutrition Education During Pregnancy on Iron-Folic Acid Supplementation Compliance and Anemia in Low- and Middle-Income Countries: A Systematic Review and Meta-analysis. *Nutrition Reviews*, 83(7), e1472–e1487. <https://doi.org/10.1093/nutrit/nuae170>
- 10 Fang, W., Reddy, O. S., & Lai, W.-F. (2024). Nutritional awareness of pregnant women and the underlying influencing factors. *Nutrition Reviews*, 82(4), 561–569. <https://doi.org/10.1093/nutrit/nuad074>
- 10 Fernández-Gómez, E., Luque-Vara, T., Moya-Fernández, P. J., López-Olivares, M., Gallardo-Vigil, M. Á., & Enrique-Mirón, C. (2020). Factors Influencing Dietary Patterns during Pregnancy in a Culturally Diverse Society. *Nutrients*, 12(11). <https://doi.org/10.3390/nu12113242>
- 11 Hagger, M. S., & Hamilton, K. (2025). Progress on theory of planned behavior research: advances in research synthesis and agenda for future research. *Journal of Behavioral Medicine*, 48(1), 43–56. <https://doi.org/10.1007/s10865-024-00545-8>
- 5 Hanani, Z., Suyatno, & P, S. F. (2016). Faktor-faktor yang Mempengaruhi Konsumsi Sayur dan Buah pada Ibu Hamil di Indonesia (Berdasarkan Data Riskesdas 2013). *Jurnal Kesehatan Masyarakat*, 4(1), 257–266.
- 18 Iwanowicz-Palus, G., Zarajczyk, M., Bień, A., Korzyńska-Piętas, M., Krysa, J., Rahnama-Hezavah, M., & Wdowiak, A. (2022). The Relationship between Social Support, Self-Efficacy and Characteristics of Women with Diabetes during Pregnancy. In *International Journal of Environmental Research and Public Health* (Vol. 19, Issue 1, p. 304). <https://doi.org/10.3390/ijerph19010304>
- 12 Ju, I., Yel, E., & Song, H. (2025). Exploring the Cognitive, Affective, and Conative Factors of Bystander Intervention to Improve Protective Measures During Health Emergencies. *Health Communication*, 40(10), 2099–2109. <https://doi.org/10.1080/10410236.2024.2442560>
- 27 Kabahenda, M. K., & Stoecker, B. J. (2024). Associations between maternal dietary intake and nutritional status with fetal growth at 14 to 26 weeks gestation: a cross-sectional study. *BMC Nutrition*, 10(1), 77. <https://doi.org/10.1186/s40795-024-00885-3>
- 22 Kim, Sehee. (2025). Validity and Reliability of the Korean Version of the Self-Efficacy Scale for Healthy Eating and Physical Activity Among Pregnant Women. *Western Journal of Nursing Research*, 47(9), 887–896. <https://doi.org/10.1177/01939459251351464>
- 13 Lim, E. J., Ong, C., Said Abdul Rashid, N. S., Lee, J. J., Chew, J., & Chua, M. C. (2025). Food

Author names dkk / Scientific Periodical of Public Health and Coastal 4(2),2022 , halaman 87-99 (10pt, all caps, Cambria, Normal)

- Accessibility and Nutritional Outcomes Among Food-Insecure Pregnant Women in Singapore. In *Nutrients* (Vol. 17, Issue 5, p. 835). <https://doi.org/10.3390/nu17050835>
- Monterrosa, Eva C, Frongillo, Edward A, Drewnowski, Adam, de Pee, Saskia, & Vandevijvere, Stefanie. (2020). Sociocultural Influences on Food Choices and Implications for Sustainable Healthy Diets. *Food and Nutrition Bulletin*, 41(2_suppl), 59S-73S. <https://doi.org/10.1177/0379572120975874>
- Mulyati, S., Mahanani, D., Kurniawan, Y. F., Ayu, I. Y., Budianto, E. T., Sekundaria, R. D., Fakhirah, A., & Fatimah, G. N. (2025). Hubungan perilaku pola makan pada status gizi ibu hamil untuk pencegahan resiko stunting di Puskesmas Kemayoran Jakarta Pusat tahun 2025. *Indonesian Journal of Health Science*, 5(5), 1177–1184. <https://doi.org/10.54957/ijhs.v5i5.1755>
- Nabillah, D., Putri, S. E., Krairunnas, Mulyani, I., & Lubis, A. I. (2024). The Relationship between Consumption Patterns and Nutritional Status of Pregnant Women at the Kuta Padang Layung Health Center. *JNS: Journal of Nutrition Science*, 5(2), 4–8. <https://doi.org/10.35308/jns.v5i2.10812>
- Nasution, A., & Nasution, A. S. (2020). Nutrition Puzzles as an Effort to Promote Balanced Nutrition Behavior in Students. *Media Kesehatan Masyarakat Indonesia*, 16(1), 89–99. <https://doi.org/10.30597/mkmi.v16i1.8606>
- Nasution, A. S., Fernandez, J. C., Putri, R., & Dwipa, U. (2024). Implementasi Program Nutrition Goes To School Bagi Pesantren Kabupaten Bogor. *Jupemi*, 2, 27–31.
- Nilawati, W., Ismawati, R., & Monalisa, D. (2024). The Influence of Video-Based Nutrition Education on Knowledge of Pregnancy Nutrition in Pregnant Women in Sinar Pasemah Village. *Journal of Educational Innovation and Public Health*, 3(1), 01–11. <https://doi.org/10.55606/innovation.v3i1.3295>
- Olloqui-Mundet, M. J., Cavia, M. D., Alonso-Torre, S. R., & Carrillo, C. (2024). Dietary Habits and Nutritional Knowledge of Pregnant Women: The Importance of Nutrition Education. In *Foods* (Vol. 13, Issue 19, p. 3189). <https://doi.org/10.3390/foods13193189>
- Rockliffe, L., Peters, S., Heazell, A. E. P., & Smith, D. M. (2021). Factors influencing health behaviour change during pregnancy: a systematic review and meta-synthesis. *Health Psychology Review*, 15(4), 613–632. <https://doi.org/10.1080/17437199.2021.1938632>
- Schwarzer, R., & Hamilton, K. (2020). Changing Behavior Using the Health Action Process Approach. *The Handbook of Behavior Change*, 89–103. <https://doi.org/10.1017/9781108677318.007>
- Suryani, D., Krisnasary, A., Pratiwi, B. A., & Yandrizal, Y. (2024). The Influence of Maternal Knowledge Increase about Pregnant Women's Nutrition on Pregnancy Outcomes in Bengkulu City. *Amerta Nutrition*, 8(3SP), 17–23. <https://doi.org/10.20473/amnt.v8i3SP.2024.17-23>
- Yu, M., Chen, H., & Li, N. (2024). Influence of nutritional guidance during pregnancy on nutritional status and pregnancy outcome of pregnant women. *African Journal of Reproductive Health*, 28(12), 29–37. <https://doi.org/10.29063/ajrh2024/v28i12.3>