

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/226722512>

Green–Naghdi theory, part A: Green–Naghdi (GN) equations for shallow water waves

Article *in* Journal of Marine Science and Application · September 2011
DOI: 10.1007/s11804-011-1066-1

CITATIONS
39

3 authors:



William Webster
University of California, Berkeley
18 PUBLICATIONS 683 CITATIONS
[SEE PROFILE](#)



Binbin Zhao
Harbin Engineering University
64 PUBLICATIONS 690 CITATIONS
[SEE PROFILE](#)

READS
1,817



Wenyang Duan
Harbin Engineering University
103 PUBLICATIONS 1,139 CITATIONS
[SEE PROFILE](#)



ISSN 1671-9433

JOURNAL OF MARINE SCIENCE AND APPLICATION

船舶与海洋工程学报

Vol. 10 No. 3 Sep. 2011



CSNAME



HEU



Springer

CONTENTS

RESEARCH PAPERS

- Green-Naghdi Theory, Part A: Green-Naghdi (GN) Equations for Shallow Water Waves
.....William C. Webster, Wenyang Duan and Binbin Zhao (253)
- A Free Surface Frequency Domain Green Function with Viscous Dissipation and Partial Reflections from Side Walls
.....Hongde Qin, Jing Shen and Xiaobo Chen (259)
- Numerical Simulation of Nonlinear Wave Force on a Quasi-ellipse Caisson
.....Yongxue Wang, Xiaozhong Ren and Guoyu Wang (265)
- A New Rational-based Optimal Design Strategy of Ship Structure Based on Multi-level Analysis and Super-element Modeling Method
.....Li Sun and Deyu Wang (272)
- Experimental Investigation of TARMAX Model for Modeling of Hydrodynamic Forces on Cylinder-like Structures
.....M. A. Lotfollahi Yaghin, A. Mojtahedi, M. M. Etefagh and M. H. Aminfar (281)
- Numerical Analysis of the High Skew Propeller of an Underwater Vehicle
.....Hassan Ghasseni and Parviz Ghadimi (289)
- Research on the Fragment Impact Resistance of a Composite Mast
.....Wei Zhang, Chao Wu, Wenshan Yang, Shanshan Xu and Shaofei Ren (300)
- Hydrodynamic Comparison of a Semi-submersible, TLP, and Spar: Numerical Study in the South China Sea Environment
.....Binbin Li, Kun Liu, Gongwei Yan and Jinping Ou (306)
- Safety Assessment for a Side-by-side Offloading Mooring System
.....Liping Sun, Qiang He and Shangmao Ai (315)
- Finite Element Simplified Fatigue Analysis Method for a Non-tubular Joint of an Offshore Jacket Platform
.....Qinghua Bao and Heng Feng (321)
- Global Robust and Adaptive Output Feedback Dynamic Positioning of Surface Ships
.....Khac Duc Do (325)
- Design of an Adaptive Controller for Dive-plane Control of a Torpedo-shaped AUV
.....Jian Cao, Yumin Su and Jinxin Zhao (333)
- Research on a Multi-grid Model for Passenger Evacuation in Ships
.....Miao Chen, Duanfeng Han and Haipeng Zhang (340)
- A Model of Application System for Man-Machine-Environment System Engineering in Vessels Based on IDEF0
.....Zhen Shang, Changhua Qiu and Shifan Zhu (347)
- Research on the Strategy of Underwater United Detection Fusion and Communication Using Multi-sensor
.....Zhenhua Xu, Jianguo Huang, Hai Huang and Qunfei Zhang (358)
- Experimental and Numerical Procedures of a Sonar Platform with a Sound Absorption Wedge
.....Danzhu Yu, Xiongliang Yao and Shaoshi Dai (364)
- Underwater Simultaneous Localization and Mapping Based on Forward-looking Sonar
.....Tiedong Zhang, Wenjing Zeng and Lei Wan (371)

TRENDS

- 9th International Conference on Advances in Fluid Mechanics (AFM 2012).....(280)
- 31th International Conference on Ocean, Offshore and Arctic Engineering (OMAE 2012).....(370)

Green-Naghdi Theory, Part A: Green-Naghdi (GN) Equations for Shallow Water Waves

William C. Webster¹, Wenyang Duan^{2*} and Binbin Zhao²

1. Department of Civil & Environmental Engineering, University of California, Berkeley CA 94720-1710, USA

2. College of Shipbuilding Engineering, Harbin Engineering University, 150001 Harbin, China

Abstract: In this work, Green-Naghdi (GN) equations with general weight functions were derived in a simple way. A wave-absorbing beach was also considered in the general GN equations. A numerical solution for a level higher than 4 was not feasible in the past with the original GN equations. The GN equations for shallow water waves were simplified here, which make the application of high level (higher than 4) equations feasible. The linear dispersion relationships of the first seven levels were presented. The accuracy of dispersion relationships increased as the level increased. Level 7 GN equations are capable of simulating waves out to wave number times depth $kd < 26$. Numerical simulation of nonlinear water waves was performed by use of Level 5 and 7 GN equations, which will be presented in the next paper.

Keywords: Green-Naghdi (GN) equations; dispersion relation; wave-absorbing beach; shallow-water waves

Article ID: 1671-9433(2011)03-0253-06

1 Introduction

The present work focuses on the simplification of Green-Naghdi (GN) equations. The linear dispersion relationships of the first seven levels of the GN equations are also obtained. The Green-Naghdi equations were originally developed by Green and Naghdi in 1974 to analyze some nonlinear free-surface flows. After the successful application of the method to nonlinear ship wave-making problems (Ertekin *et al.*, 1986), the method was applied to many nonlinear water wave problems. Later, the model has been extended to deep-water waves by Webster and Kim (1990) and Xu *et al.* (1997) in two and three dimensions, respectively. Demirebilek and Webster (1992, 1999) applied the GN theory to shallow water and nonlinear wave propagation problems in two dimensions. Ertekin and Kim (1999) studied the hydroelastic behavior of a three-dimensional mat-type very large floating structures (VLFS) in oblique waves by use of the linear GN theory. Xia *et al.* (2008) studied the two-dimensional, nonlinear hydroelasticity of a mat-type very large floating structure (VLFS) within the scope of linear beam theory for structures and the nonlinear Level 1 GN theory for the fluids Zhao *et al.* (2011) applied the Level 2 GN equations to tsunami simulation.

The original GN governing equations were not limited to irrotational flows. Kim *et al.* (2001) derived irrotational Green-Naghdi (IGN) equations. The deep-water version of the IGN equations has been derived and applied successfully to simulate ocean waves of extreme heights (Kim and Ertekin, 2000). The IGN equations were also derived for finite water-depth conditions, and tested numerically to show their self-convergence and accuracy (Kim *et al.*, 2003).

Previously, the numerical solution for a level higher than 4 was not feasible with original GN equations. Zhao and Duan (2010) performed numerical calculations for the GN equations up to Level 3. They found that the results of Level 3 GN equations are better than that of Level 2 GN equations when compared with the experimental values. They did not perform calculations for a level higher than 4 because of the complexity of the original GN equations. As will be shown below, the numerical results of Level 5 GN equations are much better than Level 3 GN equations.

The simplification of the original GN equations is necessary to facilitate the application of the equations with a level higher than 4. Derivation of the original GN equations (Demirebilek and Webster, 1992) is complex and this method can be improved. In each numerical test, the self-convergence of the GN equations should be performed first. Then, the converged solutions can be compared with known experimental values and/or other solutions.

This paper is organized as follows. In Section 2 the GN equations with a wave-absorbing beach are derived in a

Received date: 2011-07-04.

Foundation item: Supported by the Special Fund for Basic Scientific Research of Central Colleges, Harbin Engineering University (Harbin), the National Natural Science Foundation of China, Doctor Subject Foundation of the Ministry of Education of China, and the "111" project (B07019).

***Corresponding author Email:** duanwenyang@hrbeu.edu.cn

© Harbin Engineering University and Springer-Verlag Berlin Heidelberg 2011

general case. Section 3 provides the GN equations for shallow water waves, which are much simpler than the original GN equations. Section 4 provides the linear dispersion relationships of the first seven levels of GN equations for shallow water waves. Some conclusions are given in Section 5.

2 GN equations with general weight functions

The free-surface flow of an inviscid and incompressible fluid in water of variable depth is considered. The coordinate system is chosen such that the z -axis directs against gravity and the Oxy -plane is the still water level. The location of the free surface is denoted by $z = \beta(x, y, t)$ and the bottom as $z = \alpha(x, y, t)$. It should be noted that the bottom is allowed to vary with time in the GN model.

The conservation equations of momentum of the original GN model (Demirbilek and Webster, 1992) start from

$$\rho \bar{v}_{,i} + (\rho v_i \bar{v})_{,i} = -p_{,i} \bar{e}_i - \rho g \bar{e}_3 \quad (1)$$

Here we derive the conservation equations of momentum of the GN model from

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} + v \frac{\partial u}{\partial y} + w \frac{\partial u}{\partial z} + \mu u = -\frac{1}{\rho} \frac{\partial p}{\partial x} \quad (2a)$$

$$\frac{\partial v}{\partial t} + u \frac{\partial v}{\partial x} + v \frac{\partial v}{\partial y} + w \frac{\partial v}{\partial z} + \mu v = -\frac{1}{\rho} \frac{\partial p}{\partial y} \quad (2b)$$

$$\frac{\partial w}{\partial t} + u \frac{\partial w}{\partial x} + v \frac{\partial w}{\partial y} + w \frac{\partial w}{\partial z} + \mu w = -\frac{1}{\rho} \left(\frac{\partial p}{\partial z} + \rho g \right) \quad (2c)$$

where μ , a given function of (x, y) , is the artificial damping coefficient. The terms μu , μv , and μw represent the wave-absorbing beach. Xu *et al.* (2007) introduced these terms into the GN equations for deep water waves. Although Eq.(2) and Eq.(1) are the same in essence, Eq.(1) makes the derivation of GN equations harder. The other conditions which the GN model should observe are:

Continuity condition:

$$\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} = 0 \quad (3)$$

Kinematic free surface and bottom condition:

$$w - \frac{\partial \beta}{\partial t} - u \frac{\partial \beta}{\partial x} - v \frac{\partial \beta}{\partial y} = 0 \quad (z = \beta) \quad (4)$$

$$w - \frac{\partial \alpha}{\partial t} - u \frac{\partial \alpha}{\partial x} - v \frac{\partial \alpha}{\partial y} = 0 \quad (z = \alpha) \quad (5)$$

Dynamic free surface and bottom condition:

$$p|_{z=\beta} = \hat{p}(x, y, t) \quad (6)$$

$$p|_{z=\alpha} = \bar{p}(x, y, t) \quad (7)$$

In the GN model, the velocity field is expressed approximately as

$$\begin{cases} u(x, y, z, t) = \sum_{n=0}^K u_n(x, y, t) \lambda_n(z) \\ v(x, y, z, t) = \sum_{n=0}^K v_n(x, y, t) \lambda_n(z) \\ w(x, y, z, t) = \sum_{n=0}^K w_n(x, y, t) \lambda_n(z) \end{cases} \quad (8)$$

where $\lambda_n(z)$ are called shape functions. The shape functions should accurately represent the vertical structure of the flow field being dealt with. For shallow water problems, $1, z, z^2, \dots$, can be chosen as shape functions. For deep water waves, $e^{az}, e^{az^2}, e^{az^3}, \dots$, can be chosen as shape functions. The parameter a is a representative wave number. The coefficients u_n, v_n and w_n are unknown functions. K is called the level of the Green-Naghdi theory.

By use of Eq.(8), the equations of GN model can easily be obtained. They are:

Continuity condition:

$$\sum_{n=0}^K \left(\frac{\partial u_n}{\partial x} + \frac{\partial v_n}{\partial y} \right) \lambda_n(z) + \sum_{n=0}^K w_n \frac{d\lambda_n(z)}{dz} = 0 \quad (9)$$

Kinematic free surface and bottom condition:

$$\frac{\partial \beta}{\partial t} = \sum_{n=0}^K \lambda_n(\beta) \left(w_n - \frac{\partial \beta}{\partial x} u_n - \frac{\partial \beta}{\partial y} v_n \right) \quad (10)$$

$$\frac{\partial \alpha}{\partial t} = \sum_{n=0}^K \lambda_n(\alpha) \left(w_n - \frac{\partial \alpha}{\partial x} u_n - \frac{\partial \alpha}{\partial y} v_n \right) \quad (11)$$

A weak formulation similar to the Galerkin approach is employed where the shape functions λ_n are used as weighting functions to develop K approximate equations which express the conservation of momentum in some integral sense.

Taking Eq.(2a) for example, the following can easily be gotten:

$$\sum_{m=0}^K \left[\frac{\partial u_m}{\partial t} \lambda_m + \sum_{r=0}^K \left(\frac{\partial u_m}{\partial x} u_r + \frac{\partial u_m}{\partial y} v_r \right) \lambda_r \lambda_m + \sum_{r=0}^K w_r \lambda_r \frac{d\lambda_m}{dz} \lambda_m + \mu u_m \lambda_m \right] = -\frac{1}{\rho} \frac{\partial p}{\partial x} \quad (12)$$

Multiply (12) by each λ_n and integrating through the vertical direction results from bottom α to free surface β . The result is:

$$\sum_{m=0}^K \left[\frac{\partial u_m}{\partial t} S_m + \sum_{r=0}^K \left(\frac{\partial u_m}{\partial x} u_r + \frac{\partial u_m}{\partial y} v_r \right) S_{mr} + \sum_{r=0}^K u_m w_r S_{mr} + \mu u_m S_m \right] = \frac{1}{\rho} \left[-\frac{\partial P_n}{\partial x} + \hat{p} \lambda_n(\beta) \frac{\partial \beta}{\partial x} - \bar{p} \lambda_n(\alpha) \frac{\partial \alpha}{\partial x} \right] \quad (13)$$

where

$$S_m = \int_{\alpha}^{\beta} \lambda_m \lambda_n dz, \quad S_{mr} = \int_{\alpha}^{\beta} \lambda_m \lambda_r \lambda_n dz, \quad S_m^* = \int_{\alpha}^{\beta} \frac{d\lambda_m}{dz} \lambda_r \lambda_n dz, \quad P_n = \int_{\alpha}^{\beta} p \lambda_n dz.$$

Similarly, the other two conservation equations of momentum can be obtained from Eqs.(2b) and (2c) easily. They are

$$\sum_{m=0}^K \left[\frac{\partial v_m}{\partial t} S_m + \sum_{r=0}^K \left(\frac{\partial v_m}{\partial x} u_r + \frac{\partial v_m}{\partial y} v_r \right) S_{mr} + \sum_{r=0}^K v_m w_r S_{mr} + \mu v_m S_m \right] = \frac{1}{\rho} \left[-\frac{\partial P_n}{\partial y} + \hat{p} \lambda_n(\beta) \frac{\partial \beta}{\partial y} - \bar{p} \lambda_n(\alpha) \frac{\partial \alpha}{\partial y} \right] \quad (14)$$

$$\sum_{m=0}^K \left[\frac{\partial w_m}{\partial t} S_m + \sum_{r=0}^K \left(\frac{\partial w_m}{\partial x} u_r + \frac{\partial w_m}{\partial y} v_r \right) S_{mr} + \sum_{r=0}^K w_m w_r S_{mr} + \mu w_m S_m \right] = \frac{1}{\rho} \left[P_n^* - \rho g S_n - \hat{p} \lambda_n(\beta) + \bar{p} \lambda_n(\alpha) \right] \quad (15)$$

for $n = 0, 1, 2, \dots, K$. where $S_n = \int_{\alpha}^{\beta} \lambda_n dz$, $P_n^* = \int_{\alpha}^{\beta} p \frac{d\lambda_n}{dz} dz$.

Eqs.(6), (7), (9), (10), (11), (13), (14), and (15) form the GN equations with general weight functions.

3 GN equations for shallow water waves

For shallow water problems, the polynomial weighting function set is chosen as:

$$\lambda_n(z) = z^n \quad (16)$$

The velocity field is given by

$$\begin{cases} u(x, y, z, t) = u_0 + u_1 z + u_2 z^2 + \dots + u_K z^K \\ v(x, y, z, t) = v_0 + v_1 z + v_2 z^2 + \dots + v_K z^K \\ w(x, y, z, t) = w_0 + w_1 z + w_2 z^2 + \dots + w_K z^K \end{cases} \quad (17)$$

The equations for kinematic boundary conditions, conservation of mass, and momentum can now be reduced using Eq.(16). The kinematic boundary conditions becomes

$$\frac{\partial \beta}{\partial t} = \sum_{n=0}^K \beta^n \left(w_n - \frac{\partial \beta}{\partial x} u_n - \frac{\partial \beta}{\partial y} v_n \right) \quad (18)$$

$$\frac{\partial \alpha}{\partial t} = \sum_{n=0}^K \alpha^n \left(w_n - \frac{\partial \alpha}{\partial x} u_n - \frac{\partial \alpha}{\partial y} v_n \right) \quad (19)$$

The continuity equation becomes

$$\left(\frac{\partial u_K}{\partial x} + \frac{\partial v_K}{\partial y} \right) z^K + \sum_{n=0}^{K-1} \left(\frac{\partial u_n}{\partial x} + \frac{\partial v_n}{\partial y} + w_{n+1} (n+1) \right) z^n = 0 \quad (20)$$

If Eq.(20) is to be held everywhere, each coefficient of z^n must be set to zero, that is

$$\frac{\partial u_K}{\partial x} + \frac{\partial v_K}{\partial y} = 0 \quad (21)$$

$$\frac{\partial u_n}{\partial x} + \frac{\partial v_n}{\partial y} + (n+1) w_{n+1} = 0 \quad (n = 0, 1, \dots, K-1) \quad (22)$$

Demirbilek and Webster (1992) introduced the restricted GN theory. They restricted the last component of the director. That is

$$u_K = 0 \quad (23)$$

$$v_K = 0 \quad (24)$$

The conditions for conservation of momentum become

$$E_n = \frac{1}{\rho} \left[-\frac{\partial P_n}{\partial x} + \hat{p} \beta^n \frac{\partial \beta}{\partial x} - \bar{p} \alpha^n \frac{\partial \alpha}{\partial x} \right] \quad (25)$$

$$F_n = \frac{1}{\rho} \left[-\frac{\partial P_n}{\partial y} + \hat{p} \beta^n \frac{\partial \beta}{\partial y} - \bar{p} \alpha^n \frac{\partial \alpha}{\partial y} \right] \quad (26)$$

$$G_n = \frac{1}{\rho} \left[P_n^* - \rho g S_n - \hat{p} \beta^n + \bar{p} \alpha^n \right] \quad (27)$$

For $n = 0, 1, 2, \dots, K$. Where

$$E_n = \sum_{m=0}^K \left[\frac{\partial u_m}{\partial t} S_m + \sum_{r=0}^K \left(\frac{\partial u_m}{\partial x} u_r + \frac{\partial u_m}{\partial y} v_r \right) S_{mr} + \sum_{r=0}^K u_m w_r S_{mr} + \mu u_m S_m \right]$$

$$F_n = \sum_{m=0}^K \left[\frac{\partial v_m}{\partial t} S_m + \sum_{r=0}^K \left(\frac{\partial v_m}{\partial x} u_r + \frac{\partial v_m}{\partial y} v_r \right) S_{mr} + \sum_{r=0}^K v_m w_r S_{mr} + \mu v_m S_m \right]$$

$$G_n = \sum_{m=0}^K \left[\frac{\partial w_m}{\partial t} S_m + \sum_{r=0}^K \left(\frac{\partial w_m}{\partial x} u_r + \frac{\partial w_m}{\partial y} v_r \right) S_{mr} + \sum_{r=0}^K w_m w_r S_{mr} + \mu w_m S_m \right]$$

$$S_n = \int_{\alpha}^{\beta} z^n dz, \quad S_m = \int_{\alpha}^{\beta} z^{m+n} dz, \quad S_{mr} = \int_{\alpha}^{\beta} z^{m+r+n} dz, \quad S_m^* = m \int_{\alpha}^{\beta} z^{m+n-1} dz$$

Notice that

$$P_n^* = \int_{\alpha}^{\beta} p \frac{d\lambda_n}{dz} dz = n \int_{\alpha}^{\beta} p z^{n-1} dz = n P_{n-1} \quad (28)$$

With $n = 0$, Eq.(27) becomes

$$G_0 = \frac{1}{\rho} (-\rho g S_0 - \hat{p} + \bar{p}) \quad (29)$$

One can easily get the pressure on the bottom from Eq.(29); it can be expressed by

$$\bar{p} = \rho G_0 + \rho g S_0 + \hat{p} \quad (30)$$

Eqs.(25) and (26) can be rewritten as

$$\frac{\partial P_n}{\partial x} = -\rho E_n + \hat{p} \beta^n \frac{\partial \beta}{\partial x} - \bar{p} \alpha^n \frac{\partial \alpha}{\partial x} \quad (31)$$

$$\frac{\partial P_n}{\partial y} = -\rho F_n + \hat{p} \beta^n \frac{\partial \beta}{\partial y} - \bar{p} \alpha^n \frac{\partial \alpha}{\partial y} \quad (32)$$

By using of Eq.(28), one can rewrite Eq.(27) as

$$P_n = \frac{1}{(n+1)} (\rho G_{n+1} + \rho g S_{n+1} + \hat{p} \beta^{n+1} - \bar{p} \alpha^{n+1}) \quad (33)$$

$\partial P_n / \partial x$ and $\partial P_n / \partial y$ are eliminated from Eqs.(31), (32), and (33). The following equations are obtained:

$$\frac{\partial}{\partial x} (\rho G_{n+1} + \rho g S_{n+1} + \hat{p} \beta^{n+1} - \bar{p} \alpha^{n+1}) - \quad (34)$$

$$(n+1) \left(-\rho E_n + \hat{p} \beta^n \frac{\partial \beta}{\partial x} - \bar{p} \alpha^n \frac{\partial \alpha}{\partial x} \right) = 0$$

$$\frac{\partial}{\partial y} (\rho G_{n+1} + \rho g S_{n+1} + \hat{p} \beta^{n+1} - \bar{p} \alpha^{n+1}) - \quad (35)$$

$$(n+1) \left(-\rho F_n + \hat{p} \beta^n \frac{\partial \beta}{\partial y} - \bar{p} \alpha^n \frac{\partial \alpha}{\partial y} \right) = 0$$

Eqs.(34) and (35) can be simplified to

$$\frac{\partial}{\partial x} (G_{n+1} + g S_{n+1}) + (n+1) E_n + \quad (36)$$

$$\beta^{n+1} \frac{\partial}{\partial x} \left(\frac{\hat{p}}{\rho} \right) - \alpha^{n+1} \frac{\partial}{\partial x} \left(\frac{\bar{p}}{\rho} \right) = 0$$

$$\frac{\partial}{\partial y} (G_{n+1} + g S_{n+1}) + (n+1) F_n + \quad (37)$$

$$\beta^{n+1} \frac{\partial}{\partial y} \left(\frac{\hat{p}}{\rho} \right) - \alpha^{n+1} \frac{\partial}{\partial y} \left(\frac{\bar{p}}{\rho} \right) = 0$$

for $n = 0, 1, 2, \dots$

Finally, the GN equations for shallow water waves can be gotten. They are

$$\frac{\partial \beta}{\partial t} = \sum_{n=0}^K \beta^n \left(w_n - \frac{\partial \beta}{\partial x} u_n - \frac{\partial \beta}{\partial y} v_n \right) \quad (38)$$

$$\frac{\partial}{\partial x} (G_n + g S_n) + n E_{n-1} - \alpha^n \frac{\partial}{\partial x} (G_0 + g S_0) + (\beta^n - \alpha^n) \frac{\partial}{\partial x} \left(\frac{\hat{p}}{\rho} \right) = 0 \quad (39)$$

$$\frac{\partial}{\partial y} (G_n + g S_n) + n F_{n-1} - \alpha^n \frac{\partial}{\partial y} (G_0 + g S_0) + (\beta^n - \alpha^n) \frac{\partial}{\partial y} \left(\frac{\hat{p}}{\rho} \right) = 0 \quad (40)$$

for $n = 1, 2, 3, \dots, K$. Where

$$E_n = \sum_{m=0}^K \left[\frac{\partial u_m}{\partial t} S_{mn} + \sum_{r=0}^K \left(\frac{\partial u_m}{\partial x} u_r + \frac{\partial u_m}{\partial y} v_r \right) S_{mrn} + \sum_{r=0}^K u_m w_r S_{rm}^m + \mu u_m S_{mn} \right]$$

$$F_n = \sum_{m=0}^K \left[\frac{\partial v_m}{\partial t} S_{mn} + \sum_{r=0}^K \left(\frac{\partial v_m}{\partial x} u_r + \frac{\partial v_m}{\partial y} v_r \right) S_{mrn} + \sum_{r=0}^K v_m w_r S_{rm}^m + \mu v_m S_{mn} \right]$$

$$G_n = \sum_{m=0}^K \left[\frac{\partial w_m}{\partial t} S_{mn} + \sum_{r=0}^K \left(\frac{\partial w_m}{\partial x} u_r + \frac{\partial w_m}{\partial y} v_r \right) S_{mrn} + \sum_{r=0}^K w_m w_r S_{rm}^m + \mu w_m S_{mn} \right]$$

$$S_n = \int_{\bar{x}}^{\beta} z^n dz, S_{nn} = \int_{\bar{x}}^{\beta} z^{n+n} dz, S_{mm} = \int_{\bar{x}}^{\beta} z^{m+m} dz, S_{nn}^m = m \int_{\bar{x}}^{\beta} z^{m+n-1} dz$$

$$u_K = 0, v_K = 0$$

$$w_n = -\frac{1}{n} \left(\frac{\partial u_{n-1}}{\partial x} + \frac{\partial v_{n-1}}{\partial y} \right) \text{ for } n = 1, 2, \dots, K$$

$$w_0 = \frac{\partial \alpha}{\partial t} - \sum_{n=1}^K \alpha^n \left(w_n - \frac{\partial \alpha}{\partial x} u_n - \frac{\partial \alpha}{\partial y} v_n \right) + \frac{\partial \alpha}{\partial x} u_0 + \frac{\partial \alpha}{\partial y} v_0$$

Usually, the pressure is set at free surface $\hat{p} = 0$. For each choice of K , a complete closed set of equations is developed that is independent from those for a different value of K . Thus, the kinematic models form a hierarchy depending on K and increasing in complexity with K . Obviously, this hierarchy is different from a perturbation expansion. A terminology is adopted that describes the complexity of the theory, and henceforth, a particular member of these hierarchies is referred to as the “ K th level approximation”. The GN equations with $K=1, K=2, \dots$ are called Level 1, Level 2, ... GN equations, respectively.

4 Linear dispersion relationships of GN equations

The dispersion relationships for a small amplitude linear sinusoidal wave can be obtained from the linearized forms of Eqs.(38) and (39) in two dimensions.

Take Level 1 GN equations for example, set $\hat{p} = 0, \mu = 0$, and $\alpha = -d$. Water depth d is a constant. The linearized forms of Level 1 GN equations are

$$\frac{\partial \beta}{\partial t} + d \frac{\partial u_0}{\partial x} = 0 \quad (41)$$

$$3 \frac{\partial u_0}{\partial t} + 3g \frac{\partial \beta}{\partial x} - d^2 \frac{\partial^3 u_0}{\partial x^2 \partial t} = 0 \quad (42)$$

It is assumed that the free surface $\beta(x, t)$ and coefficient $u_0(x, t)$ can be expressed as

$$\beta(x, t) = A \cos(kx - \omega t) \quad (43)$$

$$u_0(x, t) = \tilde{u}_0 \cos(kx - \omega t) \quad (44)$$

Insert Eqs.(43) and (44) into Eq.(42); it is found that:

$$\tilde{u}_0 = \frac{3gA}{c(3 + d^2 k^2)} \quad (45)$$

Insert Eqs.(43), (44), and (45) into Eq.(41), and the linear dispersion relationships of Level 1 GN equations can be gotten. They are:

$$c^2 = \frac{3gd}{3 + d^2 k^2} \quad (46)$$

The analysis method of other levels of GN equations is similar. Here the non-dimensional dispersion relationships of the first seven levels of GN equations are given. They are

$$K = 1: \quad \bar{c}^2 = \frac{3\bar{d}}{3 + \bar{d}^2} = \bar{d} - \frac{\bar{d}^3}{3} + \frac{\bar{d}^5}{9} + O(\bar{d})^7 \quad (47)$$

$$K = 2 : \bar{c}^2 = \frac{24\bar{d}(10+\bar{d}^2)}{240+104\bar{d}^2+3\bar{d}^4} = \bar{d} - \frac{\bar{d}^3}{3} + \frac{19\bar{d}^5}{144} + O(\bar{d})^7 \quad (48)$$

$$K = 3 : \quad \bar{c}^2 = \frac{15\bar{d}(420+52\bar{d}^2+\bar{d}^4)}{6300+2880\bar{d}^2+135\bar{d}^4+\bar{d}^6} \\ = \bar{d} - \frac{\bar{d}^3}{3} + \frac{2\bar{d}^5}{15} - \frac{17\bar{d}^7}{315} + \frac{2893\bar{d}^9}{132300} + O(\bar{d})^{11} \quad (49)$$

$$K = 4 : \bar{c}^2 = \frac{120\bar{d}(42336+5712\bar{d}^2+154\bar{d}^4+\bar{d}^6)}{5080320+2378880\bar{d}^2+134064\bar{d}^4+1800\bar{d}^6+5\bar{d}^8} \\ = \bar{d} - \frac{\bar{d}^3}{3} + \frac{2\bar{d}^5}{15} - \frac{17\bar{d}^7}{315} + \frac{185173\bar{d}^9}{8467200} + O(\bar{d})^{11} \quad (50)$$

$$K = 5 : \\ \bar{c}^2 = \bar{d} - \frac{\bar{d}^3}{3} + \frac{2\bar{d}^5}{15} - \frac{17\bar{d}^7}{315} + \frac{62\bar{d}^9}{2835} - \frac{1382\bar{d}^{11}}{155925} + \\ \frac{1324889\bar{d}^{13}}{368831232} + O(\bar{d})^{15} \quad (51)$$

$$K = 6 : \\ \bar{c}^2 = \bar{d} - \frac{\bar{d}^3}{3} + \frac{2\bar{d}^5}{15} - \frac{17\bar{d}^7}{315} + \frac{62\bar{d}^9}{2835} - \frac{1382\bar{d}^{11}}{155925} + \\ \frac{158986681\bar{d}^{13}}{442597478400} + O(\bar{d})^{15} \quad (52)$$

$$K = 7 : \\ \bar{c}^2 = \bar{d} - \frac{\bar{d}^3}{3} + \frac{2\bar{d}^5}{15} - \frac{17\bar{d}^7}{315} + \frac{62\bar{d}^9}{2835} - \frac{1382\bar{d}^{11}}{155925} + \frac{21844\bar{d}^{13}}{6081075} - \\ \frac{929569\bar{d}^{15}}{638512875} + \frac{620626600207\bar{d}^{17}}{1051860569760000} + O(\bar{d})^{19} \quad (53)$$

where the non-dimensionalization is performed by $\bar{d} = kd$, $\bar{c} = c\sqrt{k/g}$. Only the dispersion relationships of the first four levels of GN equations are shown here. The Taylor series of the approximate dispersion relationships of the first seven levels are given to show their accuracy, and the underlined terms in the Taylor series are the terms equal to those of the exact value, which is

$$\bar{c}_{\text{exact}}^2 = \tanh(\bar{d}) \quad (54)$$

The Taylor series of Eq.(54) is

$$\bar{c}_{\text{exact}}^2 = \bar{d} - \frac{\bar{d}^3}{3} + \frac{2\bar{d}^5}{15} - \frac{17\bar{d}^7}{315} + \frac{62\bar{d}^9}{2835} - \frac{1382\bar{d}^{11}}{155925} + \\ \frac{21844\bar{d}^{13}}{6081075} - \frac{929569\bar{d}^{15}}{638512875} + \frac{6404582\bar{d}^{17}}{10854718875} + O(\bar{d})^{19} \quad (55)$$

In Fig.1, the dispersion relationships of the first seven levels of the GN equations are compared with the exact value.

The accuracy of the dispersion relationships of the GN equations increases as the level of the equations increases. With the choice of $K=7$, the relative errors of

$(c^2 - c_{\text{exact}}^2)/c_{\text{exact}}^2$ are less than 2% for $kd < 26$.




Fig.1 Dispersion relationships of the GN equations

The high order Boussinesq equation (Madsen *et al.*, 2006; Fuhrman and Madsen, 2009) is one of the best Boussinesq equations. The dispersion relationships of the high order Boussinesq model are

$$\frac{\omega^2}{gdk^2} = \frac{1 + \kappa^2 \mu_2 + \kappa^4 \mu_4 + \kappa^6 \mu_6 + \kappa^8 \mu_8}{1 + \kappa^2 \nu_2 + \kappa^4 \nu_4 + \kappa^6 \nu_6 + \kappa^8 \nu_8 + \kappa^{10} \nu_{10}} \quad (56)$$

where

$$\kappa = kd, \quad \sigma = \frac{\bar{z}}{d}, \quad \mu_2 = \frac{1}{6} - \frac{\sigma^2}{9}, \quad \mu_4 = \frac{1}{120} - \frac{\sigma^2}{54} + \frac{4\sigma^4}{567} \\ \mu_6 = \frac{\sigma^2}{270} - \frac{\sigma^3}{72} + \frac{29\sigma^4}{1701} - \frac{\sigma^5}{135} + \frac{2\sigma^6}{2835} \\ \mu_8 = \frac{\sigma^4}{7560} - \frac{\sigma^5}{1620} + \frac{17\sigma^6}{17010} - \frac{11\sigma^7}{17010} + \frac{8\sigma^8}{59535}, \quad \nu_2 = \frac{1}{2} - \frac{\sigma^2}{9} \\ \nu_4 = \frac{1}{24} - \frac{\sigma^2}{18} + \frac{4\sigma^4}{567}, \quad \nu_6 = \frac{\sigma}{120} - \frac{5\sigma^2}{216} + \frac{\sigma^3}{54} - \frac{5\sigma^4}{1134} + \frac{\sigma^5}{945} - \frac{\sigma^6}{2835} \\ \nu_8 = \frac{\sigma^3}{1080} - \frac{\sigma^4}{252} + \frac{7\sigma^5}{1215} - \frac{\sigma^6}{315} + \frac{4\sigma^7}{8505} + \frac{\sigma^8}{59535} \\ \nu_{10} = \frac{\sigma^5}{113400} - \frac{\sigma^6}{22680} + \frac{2\sigma^7}{25515} - \frac{\sigma^8}{17010} + \frac{\sigma^9}{59535} - \frac{\sigma^{10}}{893025}$$

The choice of $\sigma = 0.5$ leads to an accuracy, which is very similar to the Padé(8,10) expansion, and less than 2% errors are found for $kd < 25.7$.

The non-dimensional dispersion relationships of the high order Boussinesq equations (Madsen *et al.*, 2006) can be written as:

$$\bar{c}^2 = \frac{\bar{d}(1 + \bar{d}^2 \mu_2 + \bar{d}^4 \mu_4 + \bar{d}^6 \mu_6 + \bar{d}^8 \mu_8)}{1 + \bar{d}^2 \nu_2 + \bar{d}^4 \nu_4 + \bar{d}^6 \nu_6 + \bar{d}^8 \nu_8 + \bar{d}^{10} \nu_{10}} \quad (57)$$

where $\bar{d} = \kappa = kd$. In Fig. 2, the dispersion relationships of the Level 7 GN equations are compared with the exact value and high order Boussinesq equations.

It should be noted that there are fourth and fifth-derivatives in the high order Boussinesq equations. In contrast, the highest order derivatives in GN equations are third-derivatives no matter how high the level is.




Fig.2 Comparison of dispersion relationships between Level 7 GN equations and high order Boussinesq equations (Madsen et al., 2006)

5 Conclusions

In this paper, the GN equations with general weight functions were derived in a simple way. A wave-absorbing beach was also considered in the general GN equations. The numerical solution for a level higher than 4 was not feasible in the past with the original GN equations. After the simplification of GN equations for shallow water waves, application of high level (higher than 4) equations was feasible. Numerical calculations has been performed by use of the simplified Level 5 and Level 7 GN equations, which will be presented in the next paper. The linear dispersion relationships of the first seven levels of GN equations were also presented here. The accuracy of dispersion relationships increased as the level increased. From Eqs.(47)–(53), it is found that Level 2, 4, and 6 did not improve much in their dispersion relationships compared with previous levels. The Level 1, 2, 3, 4, 5, 6, and 7 GN equations have $O(\bar{d})^5$, $O(\bar{d})^5$, $O(\bar{d})^9$, $O(\bar{d})^9$, $O(\bar{d})^{13}$, $O(\bar{d})^{13}$, and $O(\bar{d})^{17}$ error on dispersion relationship compared with the exact value, respectively. In each numerical test, the self-convergence of the GN equations should be performed first by use of Level 1, 3, 5, and 7 GN equations. Then, the converged solutions can be compared with known experimental values and/or other solutions. Numerical simulation of nonlinear water waves will be presented in the next paper.

References


- Demirbilek Z, Webster WC (1992). Application of the Green-Naghdi theory of fluid sheets to shallow-water waves. Report 1, Model formulation. US Army Engineer Waterways Experiment Station, Coastal Engineering Research Center Technical Report No. CERC-92-11, Vicksburg, MS, 45.
- Demirbilek Z, Webster WC (1999). The Green-Naghdi theory of fluid sheets for shallow-water waves. In: Herbich JB (Ed.), *Developments in Offshore Engineering*, Gulf Pub., Houston.
- Ertekin RC, Kim JW (1999). Hydroelastic response of a mat-type structure in oblique shallow-water waves. *Journal of Ship Research*, **43**(4), 229-240.
- Ertekin RC, Webster WC, Wehausen JV (1986). Waves caused by a moving disturbance in a shallow channel of finite width. *Journal of Fluid Mechanics*, **169**, 272-292.
- Fuhrman DR, Madsen PA (2009). Tsunami generation, propagation, and run-up with a high-order Boussinesq model. *Coastal Engineering*, **56**(7), 747-758.
- Kim JW, Bai KJ, Ertekin RC, Webster WC (2001). A derivation of the Green-Naghdi equations for irrotational flows. *Journal of Engineering Mathematics*, **40**(1), 17-34.
- Kim JW, Bai KJ, Ertekin RC, Webster WC (2003). A strongly-nonlinear model for water waves in water of variable depth-the Irrotational Green-Naghdi model. *Journal of Offshore Mechanics and Arctic Engineering*, **125**, 25-32.
- Kim JW, Ertekin RC (2000). A numerical study of nonlinear wave interaction in irregular seas: irrotational Green-Naghdi model. *Marine Structure*, **13**, 331-347.
- Madsen PA, Fuhrman DR, Wang B (2006). A Boussinesq-type method for fully nonlinear waves interacting with a rapidly varying bathymetry. *Coastal Engineering*, **53**(5-6), 487-504.
- Webster WC, Kim DY (1990). The dispersion of large-amplitude gravity waves in deep water. In: *Proceedings of the 18th Symposium on Naval Hydrodynamics*, Ann Arbor, USA, 397-415.
- Xia D, Ertekin RC, Kim JW (2008). Fluid-structure interaction between a two dimensional mat-type VLFS and solitary waves by the Green-Naghdi theory. *Journal of Fluids Structures*, **24**, 527-540.
- Xu Q, Pawlowski JS, Baddour RE (1997). Development of Green-Naghdi models with a wave-absorbing beach for nonlinear, irregular wave propagation. *Journal of Marine Science and Technology*, **2**(1), 21-34.
- Zhao Binbin, Duan Wenyang, Webster WC (2011). Tsunami simulation with Green-Naghdi theory. *Ocean Engineering*, **38**(2-3), 389-396.
- Zhao Binbin, Duan Wenyang (2010). Fully Nonlinear Shallow Water Waves Simulation Using Green-Naghdi Theory. *Journal of Marine Science and Application*, **9**(1), 1-7.



William C. Webster was born in 1937. He is a professor at the University of California, Berkeley (USA). His current research interests include mooring systems nonlinear wave-body interactions, and nonlinear wave models.



Wenyang Duan was born in 1967. He is a professor and a PhD supervisor at Harbin Engineering University. His current research interests include nonlinear wave-body interactions and the SPH method.



Binbin Zhao was born in 1984. He is a lecturer at Harbin Engineering University. His current research interests include Green-Naghdi theory and mooring systems.

JOURNAL OF MARINE SCIENCE AND APPLICATION

(Quarterly, Established in June 2002)

Vol.10 No.3 September 2011

Supervised by

Ministry of Industry and Information Technology of the People's Republic of China

Sponsored by

The Chinese Society of Naval Architects and Marine Engineers

Harbin Engineering University

Edited & Published by

Editorial Department of Journal of Marine Science and Application

(1st Building 145 Nantong Street, Harbin 150001, China)

Distributed by

Harbin Post Bureau (Domestic) and Springer-Verlag (Overseas)

Tel: 86-451-82519357, 82589211

Fax: 86-451-82589211

E-mail: jmsa@hrbeu.edu.cn

Submission website: <http://mc03.manuscriptcentral.com/jmsa>

ISSN 1671-9433 CN 23-1505/T

Subscription Code: 14-323

Price: RMB ¥ 20.00

ISSN 1671-9433

